arXiv:hep-th/0004033v1 6 Apr 2000

Confinement in a Model with Condensate of the Scalar Field

V.E. Kuzmichev [a] **and V.V. Kuzmichev** [b]

Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine,
Metrolohichna St. 14b, Kiev, 03143 Ukraine

Abstract: The idea of "soft" confinement when the lifetime of hadron with respect to quark-gluon channel of decay is greater or at least of the order of some characteristic time for our Universe is considered. Within the framework of a model of three nonlinearly interacting fields the explicit form of an effective potential is found. It provides the confinement of a massive particle within the limited region of space by means of constant component of the potential which arises as a result of reorganization of vacuum of one of the scalar fields. It is shown that the lifetime of hadron being equal to the age of the Universe leads to the Higgs boson with the mass $m_H > 63.7$ GeV and realistic self-constant.

1. Introduction. The experimental data and the predictions of gauge theory of the strong interactions allow to conclude that an effective coupling constant at sufficiently small distances becomes small (phenomenon of asymptotic freedom) but at distances large in comparison with the hadron size (~ 1 fm) the coupling becomes very strong and results in quark and gluon confinement [1]. In other words the channels of decay of hadron via quarks and gluons are strictly forbidden. The nature of this prohibition is still unknown and it is introduced into the theory phenomenologically [2]. In nonrelativistic approximation a naive potential model gives a good description of the radial excitations of "hidden charm" ($c\bar{c}$) mesons if at large distances the potential is linearly or logarithmically growing to infinity but at small distances it is similar to Coulomb interaction [1]. At the distances ~ 1 fm an effective cut off of the confining potential must be realized. The bag model [3] - [7] is an example of such model where the condition that colour objects cannot appear as free particles is introduced explicitly by means of disappearance of quark current on the surface of the bag. With respect to potential model such condition is equivalent to the introduction of the unnatural infinite repulsion on the side of the external region to which quarks cannot penetrate.

In the present article we show that above mentioned cut off can be provided naturally by the condensate of the Higgs field which arises from the reorganization of the vacuum state [8] in the system of nonlinearly interacting fields. In such approach from the idea of absolute (strict) confinement (the decay of hadron into the free quarks and gluons is strictly forbidden) we come to the assumption of soft (nonstrict) confinement when the lifetime of hadron with respect to quark-gluon channel of decay is greater or at least of the order of some characteristic time for our Universe. For example, the age of our Universe can be in the role of such time.

2. Theory. We shall consider the system of three nonlinearly interacting fields ϕ, φ_1 and φ_2. The field ϕ is supposed to be massive and complex and the fields φ_i are real. For simplicity we shall not take into account their spin, colour and flavour degrees of

---

[a] e-mail: specrada@bitp.kiev.ua

[b] e-mail: specrada@bitp.kiev.ua; vvkuzmichev@yahoo.com

freedom. The corresponding generalization can be made by transition to spinor algebra. Let us define the Lorentz-invariant Lagrangian in the form

$$\mathcal{L} = \partial_\mu \phi^* \, \partial^\mu \phi - \phi^* m^2 \phi + \sum_{i=1}^{2} \left[\frac{1}{2} \, \partial_\mu \varphi_i \, \partial^\mu \varphi_i - \phi^* U_i(\varphi_i) \phi \right], \tag{1}$$

where m is the mass of the field ϕ. The functions $U_i(\varphi_i)$ are chosen in the form of anti-Higgs ($i = 1$) and Higgs ($i = 2$) potentials respectively

$$U_i(\varphi_i) = (-1)^i \left[-\frac{\mu_i^2}{2} \, \varphi_i^2 + \frac{\lambda_i}{4} \, \varphi_i^4 \right] \tag{2}$$

with constants $\mu_i^2 > 0$ and $\lambda_i > 0$. Variational principle being applied to the Lagrangian (1) leads to the set of three nonlinear equations:

$$\Box \phi = - \left[\sum_i U_i + m^2 \right] \phi, \tag{3}$$

$$\Box \varphi_i = - \left| \phi \right|^2 \, \frac{\partial U_i}{\partial \varphi_i}. \tag{4}$$

We shall limit ourselves to the consideration of the excitation spectrum of the field ϕ with energies $E' \ll 2m$. After the unitary transformation $\phi = \psi \exp(-imt)$ the equation (3) in mentioned nonrelativistic limit reduces to the Schrödinger equation

$$i \, \partial_t \, \psi = \frac{1}{2m} \left[-\nabla^2 + V \right] \psi \tag{5}$$

with the potential

$$V = \sum_i U_i(\varphi_i). \tag{6}$$

This equation describes the motion of a particle with the mass m in the field of the potential (6) being provided by the relativistic fields φ_i. We shall consider the case when a particle scatters on the potential (6) or tunnels from it. We take the axis OZ for the initial direction of its motion. It turned out that the interaction will have the form of the barrier interesting for us if we suppose that dissipation of the fields φ_i in (x, y) - plane is small. It means that the fields φ_i represent the structures stretched along the axis OZ with a cross-section $R^2 = x^2 + y^2$ small in comparison with the square of the distance z^2 at which the physical processes under consideration are observed. In this case the radial variable r in the equation (5) coincides with $|z|$ to within the terms of the order of $R^2/2\, z^2$ and fields φ_i form the local spacetime particle-like formations which are the carriers of the effective interaction along the r. Considering the amplitude $|\phi| = |\psi|$ as a slowly varying function of the variables $(x, y, z; t)$ in the above approximation we shall find the solutions of the equations (4) in the form of the solitary wave of sech-type ($i = 1$) and kink/antikink ($i = 2$) in (z, t) - plane [9, 10]. Substitution of these solutions into (6) leads to the effective potential

$$V(z, t) = \frac{\mu_1^4}{\lambda_1} \operatorname{sech}^2 (\mu_1 |\psi| s_1) \tanh^2(\mu_1 |\psi| s_1) + \frac{\mu_2^4}{4 \lambda_2} \left[\operatorname{sech}^4 \left(\frac{\mu_2}{\sqrt{2}} |\psi| s_2 \right) - 1 \right], \tag{7}$$



Figure 1: The effective potential (7) in the stationary case ($u_i = 0$) as a function of the radial variable $r = \sqrt{\xi^2 + \eta^2}$ and its section by the plane passing across the axis of symmetry.

where $s_i = (z - u_i t)/\sqrt{1 - u_i^2}$, and u_i are the velocities of the solitary waves, $-1 < u_i < 1$.

The first term in (7) is proportional to the sech-wave energy density being local. It has the shape of two humps with the maximums in the points $\mu_1 |\psi| s_1 = \pm 0.881$ and minimums at $\pm\infty$ and at the origin and organizes the interaction in the form of the barrier of the finite magnitude.

The second term in (7) describes the part of the interaction formed by the kink/antikink with the energy density localized near the value $|\psi|\, s_2 = 0$. Within limit of $|\psi|\, s_2 \to \pm\infty$ it takes the constant negative value $V_c = -\mu_2^4/4\,\lambda_2$.

Since both terms in (7) in the (z,t) - plane have a pronounced local character then the possible dependence of the amplitude $|\psi|$ on these variables does not change general form of the potential $V(z,t)$. Its concrete form depends only on the relations between the parameters μ_i, λ_i and u_i. In stationary case at $(\mu_1^4/\lambda_1) > (\mu_2^4/\lambda_2)$ it has the form of the barrier localized in the space. One of the possible variants is shown in Fig. 1.

Scattering of the particle on such potential is described by the wave function in the form of the superposition of converging and diverging waves. It is convenient to normalize it to the unit density of the incident flux, $\psi = \exp(iqz - iEt) + \psi_{scat}$, where $E = q^2/2m$ is the energy of the particle and ψ_{scat} takes into account the action of the potential. Since $\psi_{scat} \to 0$ at $r \to 0$ [11] then $|\psi| \to 1$ at $r \to \infty$. Hence at large values of $|z|$ the potential (7) tends to the constant negative value V_c which can be expressed in terms of the Higgs mass $m_H = 2\,\mu_2\,m$ and Higgs self-constant $\lambda = \lambda_2\,m^2$ corresponding to the free field φ_2,

$$V_c = -\frac{1}{64\,\lambda}\,\frac{m_H^4}{m^2}. \tag{8}$$

The appearance of the constant component (8) of the potential (7) at large distances $|z|$ can be interpreted as a consequence of the reorganization of vacuum of the field φ_2 as a result of spontaneous symmetry breaking. Its value is determined by the constant field v (condensate) originating from such reorganization and being equal to

$$v = \pm\,\frac{m_H}{2\,\sqrt{\lambda}} = \pm\,\frac{\overline{m}_H}{2\,\sqrt{\overline{\lambda}}}, \tag{9}$$

where $\overline{m}_H = m_H\,|\psi|$ and $\overline{\lambda} = \lambda\,|\psi|^2$ are the effective values of the Higgs mass and the Higgs self-constant in the presence of the field ψ (which generally speaking depend on the energy E and on the location in the spacetime).

3. Numerical example. In order to demonstrate the role of the potential V_c in the process of quantum tunneling of the particle from the region inside the barrier we shall consider the stationary case ($u_i = 0$) and replace corresponding "real" effective potential by the simpler one in the form of the sum of square steps [12]

$$V = V_1\,\theta(r - r_1)\,\theta(r_2 - r) - V_2\,\theta(r - r_2), \tag{10}$$

where V_1 and V_2 are the positive constants, r_2 is the size of the barrier, $\Delta r = r_2 - r_1$ is its width and $\theta(x)$ is the step-function.

Since we are interested only in the integral characteristics of the process then we may impose the constraints on the parameters of the potentials (7) and (10)

$$V_1 = \frac{1}{\Delta r}\int_0^{r_0} dz\overline{V(z,t)}, \qquad V_2 = -V_c, \tag{11}$$

where $\overline{V(r_0,t)} = 0$ and the bar denotes transition to the stationary case. The quantities Δr and r_1 are determined by the corresponding parameters of the hadron. Such simple model, of course, does not describe observed spectroscopy of mesons but it has two important properties: (i) there exists the domain $r < r_1$ where the quarks with the energy $E < V_1$ behave as free; (ii) at definite relation between V_1 and V_2 the quark tunneling into the range $r > r_2$ is suppressed. Indeed the direct calculations of the decay probability of the system from S - state with the energy close to the energy of the stationary state $E_0 = k_0^2/2m$ determined by the transcendent equation

$$\sqrt{V_1 - k_0^2}\,\tan k_0 r_1 = -k_0 \tag{12}$$

leads to the width for decay into quarks

$$\Gamma = \frac{8}{m}\,\frac{k_0^2\,(V_1 - k_0^2)^{3/2}\,(V_2 + k_0^2)^{1/2}}{V_1\,(V_1 + V_2)\,(1 + \sqrt{V_1 - k_0^2}\,r_1)}\,\exp\left\{-2\,\sqrt{V_1 - k_0^2}\,\Delta r\right\}. \tag{13}$$

At $V_1 \sim V_2$ the width Γ is determined mainly by the exponential factor. But at $V_2 \gg V_1$ the pre-exponential multiplier becomes essential. The width Γ decreases as $V_2^{-1/2}$ when V_2 grows and for sufficiently large value of V_2 but finite nonzero exponential factor we have $\Gamma \sim 0$.

Let us estimate V_1 and V_2 for some "realistic" values of Γ in approximation $V_1 \gg k_{0n}^2 = \pi^2 n^2/r_1^2$, where $n = 1, 2, 3 \ldots$ is the number of the S - level. We shall consider the tunneling of the quark from the ground $1S$ - state. In order to illustrate the role of V_2 at first we assume that $V_2 = 0$ and the width of the $1S$ - level equals to the decay constant of the nucleus U_{92}^{238} into α - particle and Th_{90}^{234}, $\Gamma_\alpha = 7 \times 10^{-18}\,\mathrm{s}^{-1}$. Taking the quark mass, the hadron size and the range of barrier to be equal to $m = 350$ MeV, $r_1 = 1$ fm and $\Delta r = 0.4$ fm respectively, we obtain $V_1 = 7 \times 10^2$ GeV for $\Gamma = \Gamma_\alpha$ and the energy $E_{01} = 547$ MeV for emitted quark. Since free quarks are not observed then the lifetime of hadron with respect to quark decay mode is not less then the age of the Universe with $\Gamma_0 = 2 \times 10^{-18}\,\mathrm{s}^{-1}$. For obtained V_1 the value $\Gamma = \Gamma_0$ is reached at $V_2 \approx 10^7$ GeV. The condition $E_{01} \ll V_1 \ll V_2$ is valid. If hadron decays with respect to the same quark channel during the time typical for weak interactions then for $\Gamma_w = 10^8\,\mathrm{s}^{-1}$ and $V_2 = 0$ we would have $V_1 = 95$ GeV. At such height of barrier V_1 the width $\Gamma = \Gamma_0$ is reached at $V_2 \approx 10^{55}$ GeV.

These estimations show that along with practically infinite values (such as $V_1 = \infty,\ V_2 \sim 0$, corresponding to quark bag model, or $V_1 < \infty,\ V_2 = \infty$) there also exist the values of V_1 and V_2, which provide the quark confinement while remaining finite within the energy scale of high energy physics.

Knowing V_2 we can estimate the value of admissible Higgs mass. For the value $V_2 \approx 10^7$ GeV and quark mass $m = 350$ MeV we have the following relation between the values m_H and λ,

$$m_H \approx 272\,\lambda^{1/4}\ \mathrm{GeV}. \tag{14}$$

For $\lambda > 3 \times 10^{-3}$ [13] it gives the value $m_H > 63.7$ GeV which coincides with the current theoretical and experimental estimations of the Higgs mass [14].

We should like to express our gratitude to Alexander von Humboldt Foundation (Bonn, Germany) for the assistance during the research.

References

[1] E.Leader and E.Predazzi, An Introduction to gauge theory and “New physics” (Cambridge University Press, Cambridge, 1983).

[2] W.Lucha and F.F.Schöberl, Die starke Wechselwirkung (Wissenschaftsverlag, Mannheim, Wien, Zürich, 1989).

[3] A.Chodos et al., Phys. Rev. D9 (1974) 3471; D10 (1974) 2599.

[4] T.DeGrand et al., Phys. Rev. D12 (1975) 2060.

[5] A.Martin, Phys. Lett. B67 (1977) 330.

[6] H.Gross, Phys. Lett. B68 (1977) 343.

[7] P.Hasenfrantz and J.Kuti, Phys. Rep. C75 (1978) 1.

[8] Ta-Pei Cheng and Ling-Fong Li, Gauge theory of elementary particle physics (Clarendon Press, Oxford, 1984).

[9] R.K.Bullough and P.J.Caudrey, Solitons, eds. R.K.Bullough, P.J.Caudrey (Springer-Verlag, Berlin, Heidelberg, New York, 1980).

[10] V.E.Kuzmichev and V.V.Kuzmichev, Three field dynamics in the (1+1)-dimensions (e-print, hep-th, April, 2000).

[11] R.G.Newton, Scattering theory of wave and particles (Springer-Verlag, New York, Heidelberg, Berlin, 1982).

[12] V.E.Kuzmichev and V.V.Kuzmichev, 15th International Conf. on Few-body Problems in Physics. Abstract of contributions. (Groningen, Netherland, 1997).

[13] A.D.Linde, Elementary particle physics and inflationary cosmology (Harwood, Chur, 1990).

[14] I.Hinchliffe, Eur. Phys. J. C3 (1998) 244.